



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE COMMON BROWN BATS OF PENINSULAR FLORIDA AND
SOUTHERN CALIFORNIA.

BY S. N. RHOADS.

Examination of a series of skins and skulls and alcoholic specimens of the Florida Brown Bat, in the author's collection and in the museum of the Academy of Natural Sciences of Philadelphia, shows constant racial differences from typical *Eptesicus fuscus* of Philadelphia county. These differences are similar and in the same degree and direction as those separating the two forms of Red Bat inhabiting the regions named. The Florida race may be distinguished as follows:

Eptesicus fuscus osceola subsp. nova.

Type No. 875, ad. ♂, in Coll. of S. N. Rhoads. Taken April 29, 1892, at Tarpon Springs, Fla., by W. S. Dickinson.

Description.—Similar in size and cranial characters to *fuscus*; colors deeper and darker, being of slightly varying shades of cinnamon brown as contrasted with the bistre and sepia of *fuscus*. This character is uniform in a series of eight dry skins which have never been immersed in a liquid preservative, and is peculiar to them in a comparison with a similar series of fifteen topotypes of *fuscus*.

Measurements of type, made by collector from fresh specimen: Total length 101 mm.; tail 38 mm.; hind foot $9\frac{1}{2}$ mm. Average measurements of four topotypes, 113-44-10.6.

The skull of type indicates it to be an old adult, quite as large as adult skulls of *fuscus*, but the measurements given by the collector are less than a normal average. This average corresponds closely with that of ten specimens of *fuscus* from Sing Sing, N. Y., as given in Miller's monograph of North American *Vespertilionidæ*.

Whether this subspecies is found outside the limits of peninsular Florida I am unable to state. As Miller classes the *Eptesicus* from Alabama, Georgia and Mississippi examined by him under *fuscus*, I conclude that *V. caroliniensis* of Geoffroy cannot apply to the Florida race.

In Miller's monograph, above cited, all the California *Eptesicus* examined by him are classed under *fuscus*. A specimen in the author's collection from the San Bernardino Valley, near San Bernardino, indicates a phase of coloration separating it from typical *fuscus*, in a manner such as might be reasonably predicted by any one having a knowledge of the climatic effect of this locality upon other mammals thus living on the outskirts of the Mojave Desert region. It is quite probable that Mr. Herron collected this specimen on his ranch, a few miles west of San Bernardino, or in one of his eastern trips into the edge of the desert. The exact locality is not stated on the label. The specimen may be thus characterized:

Eptesicus fuscus bernardinus subsp. nova.

Type No. 1,247, ad. ♂, Coll. of S. N. Rhoads. Taken by R. B. Herron, May 26, 1893, in the "San Bernardino Valley" (near San Bernardino), Cal.

Description.—Size and cranial characters as in *Eptesicus fuscus typicus*.

Color.—Pallid bistre above, brownish drab below; the hairs below being unicolor nearly to their roots, and those above darkening slightly only in the basilar fourth of their length. In *fuscus* the upper body hairs are darkly sooty for more than half their basilar length, the brown tips not concealing the dark under-fur. Wing membranes and ears in *bernardinus* very dark.

Measurements of type: Total length 114 mm.; tail 51 mm.; hind foot?; alar extent 304 mm. These measurements were made by the collector before skinning the specimen.

A series of four specimens from the same collector taken in the "San Bernardino Mts." in September, 1893, shows that the mountain form is inseparable from *fuscus*; one of these, however, is a perfect intergrade. *Bernardinus* differs from *peninsulæ*, O. Thomas, from southern Lower California, in its large size; *peninsulæ* appearing to be a distinct species.